

Where is the charm quark energy loss at RHIC?

Magdalena Djordjevic and Miklos Gyulassy

*Dept. Physics, Columbia University, 538 W 120-th Street,
New York, NY 10027, USA*

February 21, 2003

Abstract

Heavy quark energy loss in a hot QCD plasma is computed taking into account the competing effects due to suppression of zeroth order gluon radiation below the plasma frequency and the enhancement of gluon radiation due to first order medium induced Bremsstrahlung. The results suggest a surprising degree of cancellation between the two medium effects for charm quarks and provides a possible explanation for the null effect observed by PHENIX in the prompt electron spectrum in $Au + Au$ at 130 AGeV.

Introduction

Recent observations [1]-[8] of large suppression moderate $p_{\perp} \sim 5$ GeV hadrons produced in $Au + Au$ at the Relativistic Heavy Ion Collider (RHIC) have been interpreted as evidence for jet quenching of light quark and gluon jets [9]-[16]. Jet quenching was predicted [17, 18] to occur due to medium induced radiative energy loss of high energy partons propagating through ultra-dense QCD matter. The quenching pattern therefore provides a novel tomographic tool that can be used to map out the evolution of the plasma density. Of course, even without final state interactions, gluon radiation associated with hard QCD processes softens considerably the lowest order jet spectra. This is taken into account through the Q^2 (DGLAP) evolution of the hadronic fragmentation functions. Medium induced radiation is the extra gluon radiation that arises from higher twist final state interactions and depends on the optical thickness or opacity L/λ of the medium.

A similar quenching pattern was predicted [19, 20, 21] to occur for heavy quark (c or b) jets. However, in [22] it was pointed out that the heavy quark mass leads to a “dead cone” effect for $\theta < M/E$ that reduces the induced radiative energy loss of heavy quarks as compared to light partons. Numerical estimates indicated that the quenching of charm quarks may be approximately about a half that of light quarks. Experimentally, PHENIX data [23] on “prompt” single electron production in $Au + Au$ collisions at $\sqrt{s} = 130$ AGeV have provided a first test of heavy quark energy loss. Remarkably, no indication for a QCD medium effect was found within the admittedly large experimental errors (see also ref. [24]). In future runs, much higher statistics and wider p_T range will become accessible.

In this letter we investigate whether the apparent null effect observed for heavy quark energy loss via single electrons could be due to other medium effects that compete with the induced

radiation. Our title is motivated by a similar puzzling null effect seen previously with light quarks at much lower SPS energies [25, 26].

We study here the non-abelian analog of the Ter-Mikayelian [27]-[28] effect. In [22] the suppression of radiation below a plasma frequency cutoff was estimated to be only $\sim 10\%$ effect on the induced energy loss. However, the Ter-Mikayelian effect on the *zeroth order* in opacity $(L/\lambda)^0$ radiation has not yet been considered quantitatively. The first estimates of the influence of a plasma frequency cutoff in QCD plasmas were reported in ref. [29] using a constant plasmon mass ω_0 [30]-[32]. The k dependence of the gluon self energies and the magnitude of longitudinal radiation were not investigated. In this letter, we extend those results by taking both longitudinal as well as transverse modes consistently into account via the frequency and wavenumber dependent hard thermal (1-loop HTL) self energy $\Pi^{\mu\nu}(\omega, k)$ [33]-[37].

The dielectric properties of an isotropic plasma lead to a transverse gluon self energy $\Pi_T(\omega, \vec{k})$ with $\Pi_T(\omega_{pl}(0), 0) = \omega_{pl}^2(0) \approx \mu^2/3$, where $\mu \approx gT$ is the Debye screening mass of a plasma at temperature T in the deconfined phase. In addition, long wavelength collective longitudinal gluon modes arise with $\Pi_L(\omega_{pl}(0), 0) = \omega_{pl}^2(0)$. This dynamical gluon mass suppresses the radiation of soft $\omega < \omega_{pl}(\vec{k})$ gluons and shields the collinear $k_\perp \rightarrow 0$ singularities that arise for massless quarks.

The second part of our study is to generalize the GLV opacity expansion method [11] to compute the first order induced energy loss including the kinematic effects due to both heavy quarks and massive transverse gluons. These mass effects change the formation times of the gluons and hence the destructive (LPM) interference pattern that reduces the radiative energy loss in comparison to the incoherent Bethe-Heitler limit. For ultra-relativistic light partons, the formation times are generally long compared to nuclear dimensions. The energy loss $\Delta E^{ind} \propto L^2$ for very high energy light partons depends quadratically on the thickness of a static QCD medium [14, 10, 11], while for heavy quarks we show below that it is closer to a linear $\propto L^1$ dependence.

In this letter, we answer the question in the title by showing that there is a surprising degree of cancellation between the reduction of the zeroth order energy loss due to the QCD Ter-Mikayelian effect and the induced radiative GLV energy loss for heavy quarks. The technical details of the derivations and further discussion will be presented elsewhere [38, 39]. Our main result is that while the dead cone effect [22] is not sufficient by itself to explain the null effect by PHENIX [23], the approximate cancellation of medium effects for heavy quarks could be.

Plasmon Effect on 0^{th} order Energy Loss

Our first goal was to test the accuracy of the commonly used simplifying assumption of neglecting the $\vec{k}$ dependence of effective plasmon mass as in [9, 29, 11] and also neglecting longitudinal modes. As shown below and in more detail in [38], this assumption was found to be accurate to $\sim 10\%$ as long as the *asymptotic* plasmon mass [36], $\omega_\infty = \mu/\sqrt{2}$ rather than the long wavelength $\omega_{pl}(k=0) = \sqrt{2/3} \omega_\infty$ is employed. The accuracy of the approximation also improves dramatically as the mass of the heavy quark increases.

We computed the plasmon effect on the (zeroth order in opacity) vacuum energy energy loss, $\Delta E_{med}^{(0)} = \int dk (dI_T/dk + dI_L/dk)$ using the optical theorem in the soft gluon limit ignoring the spin of the heavy quark. The zeroth order transverse and longitudinal radiated energy

losses per wave number for an initial quark jet with four momentum $P = (E, \vec{p})$ are found to be given by

$$\begin{aligned}\frac{dI_T}{d|\vec{k}|} &= \frac{C_F}{\pi} \frac{4\vec{k}^2 \vec{p}^2 \omega_T^2 (\omega_T^2 - \vec{k}^2)}{\omega_T^2 \mu^2 - (\omega_T^2 - \vec{k}^2)^2} \int_{-1}^1 d\cos\theta \frac{\alpha_S(Q^2)}{(Q^2 - M^2)^2} \sin^2\theta \\ \frac{dI_L}{d|\vec{k}|} &= \frac{C_F}{\pi} \frac{4\vec{k}^2 \vec{p}^2 (\omega_L^2 - \vec{k}^2)}{\mu^2 - (\omega_L^2 - \vec{k}^2)} \int_{-1}^1 d\cos\theta \frac{\alpha_S(Q^2)}{(Q^2 - M^2)^2} \left(\cos\theta + \frac{\vec{k}^2}{2\vec{p}^2} \right)^2\end{aligned}\quad (1)$$

where $Q^2 = 2Pk + k^2 + M^2$ is the invariant mass of the intermediate state, $k^2 = \omega^2 - \vec{k}^2$ and $Pk = E\omega - |\vec{p}||\vec{k}|\cos\theta$ with $\omega = \omega_T(k)$ for the case of transverse radiation and $\omega = \omega_L(k)$ for longitudinal radiation. The well known dispersion relations, $\omega_T(k)$ and $\omega_L(k)$ are obtained from $\omega^2 \epsilon_T(\omega, k) - \vec{k}^2 = 0$ and $\epsilon_L(\omega, k) = 0$ respectively. Analytic expressions for ϵ_T and ϵ_L in the HTL or equivalently, semi-classical transport linear response approximations can be found in refs. [35, 36, 37]. The angular integration can be performed analytically if α_S does not run. We perform the integration with running coupling at the scale Q^2 as in conventional DGLAP evolution.




FIG 1. The reduction of the zeroth order (vacuum) energy loss for charm quark due to the QCD Ter-Mikayelian plasmon effect is shown as a function of the charm quark energy. The upper curve shows the vacuum energy loss if gluons are treated as massless and transversely polarized. The lower solid curve shows medium modified (but zeroth order in opacity) transverse fractional energy loss. The dashed curve shows the negligible additional effect of longitudinal plasmons.

The numerical results for a medium characterized by a Debye screening scale $\mu = 0.5$ GeV in Fig.1 show that the longitudinal plasmon contribution to the energy loss is indeed negligible for the range of energies considered. However, the transverse plasmon mass effect reduces the zeroth order energy loss by, $\sim 30\%$, relative to the vacuum case. The plasmon mass cutoff would effectively *enhance* the yield of high transverse momentum charm quarks were it not for the extra medium induced radiation discussed in the next section.

Medium induced radiative energy loss

The second effect of the medium on the the propagation of heavy quarks is induced radiation caused by the multiple interactions with partons in the medium. In order to compute medium induced energy loss for heavy quarks we extended the GLV method [40] to include the kinematic modification of the propagators. The new feature here is that unlike in previous applications, both the quark and and gluon plasmon mass (M and m_g respectively) are taken into account.

Since we found [38] that the dielectric plasmon effect can be well approximated numerically neglecting the longitudinal modes and using the wavenumber independent asymptotic mass $\mu/\sqrt{2}$ as the effective gluon mass, we will use this simplifying assumption in computing the induced radiative energy loss.

The heavy quark generalization of the GLV energy loss to first order in the opacity $L/\lambda = \int dx \sigma \rho$ is found to be [39]:

$$\Delta E_{ind}^{(1)} = \frac{C_F \alpha_s L}{\pi \lambda} \int_0^1 dx E \int \frac{\mu^2 dq_\perp^2}{(q_\perp^2 + \mu^2)^2} \int \frac{dk_\perp^2 \theta(xE - k_\perp)}{(\frac{4Ex}{L})^2 + (k_\perp^2 + M^2 x^2 + m_g^2)^2} \left\{ k_\perp^2 - M^2 x^2 - m_g^2 + \frac{(q_\perp^2 - k_\perp^2 + M^2 x^2 + m_g^2)(k_\perp^2 + M^2 x^2 + m_g^2)}{\sqrt{((k_\perp - q_\perp)^2 + M^2 x^2 + m_g^2)((k_\perp + q_\perp)^2 + M^2 x^2 + m_g^2)}} \right\}, \quad (2)$$

where $q_\perp$ is the magnitude of the transverse momentum transfer between a target parton and a jet, $k_\perp$ is the magnitude of the transverse momentum of the radiated gluon, and λ is the mean free path of the gluon. The simple analytic form of the destructive interference factor involving $(4Ex/L)^2$ arises for an assumed exponential distribution $\exp(-\Delta z/L)/L$ between scattering centers. The expression in $\{\dots\}$ reduces in the $M = m_g = 0$ limit to $2k_\perp^2 \theta(q_\perp - k_\perp)$, and the GLV asymptotic result eq.(127) of [40] can be recovered. We ignore the finite kinematic bounds on $q_\perp < \sqrt{6ET}$. The $q_\perp^2$ integral can then be performed analytically but is cumbersome. Qualitatively, the effect of increasing M is to reduce the relevance of the formation time factor $(4Ex/L)^2$ in the denominator of the integrand. Formally, setting this factor to zero recovers the Bethe-Heitler result with $\Delta E_{ind}^{(1)} \propto \alpha_s EL/\lambda$ modulo a logarithmic factor.

The numerical results for the first order induced radiative energy loss are shown on Fig.2 for charm and bottom quarks. We fix the effective static plasma opacity to be $L/\lambda = 4$ from the analysis of light quark quenching in $Au + Au$ at 130 GeV from the results of Levai et al [41]. We assume here that $\alpha_s = 0.3$, $\mu = 0.5$ GeV, and $\lambda = 1$ fm for the plasma parameters. In the energy range, $E \sim 5 - 15$ GeV, the Ter-Mikayelian effect reduces the induced energy loss in this extension of the GLV approach somewhat more than in the BDMS approximation[22]. However, on an absolute scale, this only corresponds to a change of $\delta(\Delta E^{(1)}/E) \sim 0.05$, which is negligible. Note that with both dead cone and plasmon mass reduction, there remains a sizeable induced energy loss fraction $\Delta E^{(1)}/E \approx 0.2$ for charm quarks while only about half that is predicted for bottom. Therefore, the dead cone effect by itself is not sufficient to explain the PHENIX heavy quark null effect.




FIG 2. The 1st order in opacity fractional energy loss for heavy quarks with (solid curves) and without (dashed curves) Ter-Mikayelian effect approximated by a constant $m_g = \mu/\sqrt{2}$ for transverse modes only. Upper curves correspond to charm, and lower to bottom quarks as a function of their energy in a plasma characterized by $\alpha_s = 0.3$, $\mu = 0.5$ GeV, and $L = 4\lambda = 4$ fm.

The net heavy quark energy loss

Fig.3 shows the competition between the medium dependence of the induced energy loss and the zeroth order energy loss taking into account the Ter-Mikayelian effect. Even in the absence of a medium ($L = 0$), a charm quark with energy ~ 10 GeV suffers an average energy loss, $\Delta E_{vac}^{(0)}/E \approx 1/3$, due to the sudden change of the color current when it is formed in the vacuum. The dielectric plasmon effect reduces this to about $\Delta E_{med}^{(0)}/E \approx 1/4$. This contribution is independent of the thickness of the plasma as long as L is not too small. For very small $L < 1/m_g$, the plasmon dispersion is smeared out due to the uncertainty principle, and $\Delta E_{med}^{(0)}$ must approach $\Delta E_{vac}^{(0)}$ from below.

The induced contribution, $\Delta E_{ind}^{(1)}$, on the other hand, increases with L . Note that for charm quarks the thickness dependence is closer to the linear Bethe-Heitler like form, L^1 , than the asymptotic energy quadratic form[10, 11]. More importantly, the two competing effects tend to cancel to a large extent for an effective thickness in the physically relevant range L .




FIG 3. The fractional energy loss for a 10 GeV charm quark is plotted versus the effective static thickness L of a plasma characterized by $\mu = 0.5$ GeV and $\lambda = 1$ fm. The dashed middle horizontal line corresponds to the energy loss in the vacuum taking into account the kinematic dead cone of radiation for heavy quarks [22]. The lower horizontal solid line shows our estimate of the reduction of the zeroth order energy loss due to the QCD analog of the Ter-Mikayelian effect. The solid curve corresponds to the net energy loss, $\Delta E_{ind}^{(1)} + \Delta E_{med}^{(0)}$.

The net energy loss, $\Delta E_{med}^{(0)} + \Delta E_{ind}^{(1)}$ is found to be remarkably close to the naive vacuum value $\Delta E_{vac}^{(0)}$ for the effective static opacity medium with $L \approx 3$ fm and $\mu \approx 0.5$ GeV, $\lambda \approx 1$ fm). These results suggest that in addition to the heavy quark dead cone effect, a high degree of cancellation between the competing medium effects may be an important factor that could explain the null effect found experimentally by PHENIX[23] for $Au + Au$ at 130 GeV. The different dependence of these effects on the plasma thickness L and on the transport properties, μ and λ , should make it possible to test this explanation by varying the beam energy, A , and centrality at RHIC and LHC energies.

In closing, we note a third mechanism that could also play a role in explaining the null effect reported by PHENIX. As shown in [42], for quark momenta below $\sim 3T$ detailed balance in a thermal medium gives rise to a positive feedback due to thermal gluon absorption that limits the energy loss. For a rapidly expanding hydrodynamic medium with transverse flow boost rapidity, η_T , even large energy loss cannot quench the spectrum below the blue shifted $p_T \sim 3Te^{\eta_T}$ thermal distribution *if* the heavy quark comes into local equilibrium. As noted in [24], for charm quarks there may in fact be only a small room between the unquenched pQCD spectrum and the fully quenched hydrodynamical spectrum in the p_T range accessible thusfar at RHIC. A decisive test of this explanation would be the observation of charm elliptic flow at RHIC. If, on the other hand, no charm elliptic flow is observed and the null effect persists to higher p_T , then the cancellation mechanism discussed above could solve the charm puzzle.

Acknowledgments: Valuable discussions with I. Vitev, Z. Lin, J. Nagle, and W. Zajc on

heavy quark production at RHIC are gratefully acknowledged. This work is supported by the Director, Office of Science, Office of High Energy and Nuclear Physics, Division of Nuclear Physics, of the U.S. Department of Energy under Grant No. DE-FG02-93ER40764.

References

- [1] K. Adcox *et al.* [PHENIX Collaboration], arXiv:nucl-ex/0207009.
- [2] C. Adler *et al.*, Phys. Rev. Lett. **89**, 202301 (2002).
- [3] S. Mioduszewski [PHENIX Collaboration], arXiv:nucl-ex/0210021.
- [4] G. J. Kunde [STAR Collaboration], arXiv:nucl-ex/0211018.
- [5] C. Adler *et al.* [STAR Collaboration], arXiv:nucl-ex/0210033.
- [6] C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **90**, 032301 (2003).
- [7] K. Filimonov [STAR Collaboration], arXiv:nucl-ex/0210027.
- [8] Proc. 15th Int. Ultra. Nucleus-Nucleus Coll. (Quark Matter 2001, Jan. SUNY), Nucl.Phys. A **698** (2002) 1c; Proc. 16th (Quark Matter 2002, July Nantes, France); Nucl. Phys. A in press.
- [9] M. Gyulassy, P. Levai, and I. Vitev, Phys. Lett. B **538**, 282 (2002) ; E. Wang and X.-N. Wang, Phys. Rev. Lett. **89**, 162301 (2002); C. A. Salgado and U. A. Wiedemann, Phys. Rev. Lett. **89**, 092303 (2002); M. Gyulassy, I. Vitev and X. N. Wang, Phys. Rev. Lett. **86**, 2537 (2001).
- [10] B.G. Zhakharov, JETP Lett. **63**, 952 (1996); R. Baier *et al.*, Nucl. Phys. B **484**, 265 (1997); U.A. Wiedemann, Nucl. Phys. B **588**, 303 (2000).
- [11] M. Gyulassy, P. Levai, and I. Vitev, Phys. Rev. Lett. **85**, 5535 (2000) ; Nucl. Phys. B **594**, 371 (2001) ; Phys. Rev. D **66**, 014005 (2002).
- [12] Gyulassy M, Wang X N, Nucl. Phys. B **420** (1994) 583; Wang X N, Gyulassy M, Plumer M, Phys. Rev. D **51** (1995) 3436
- [13] Wiedemann U A, Nucl. Phys. B **588** (2000) 303, Nucl. Phys. B **582** (2000) 409
- [14] Baier R, Dokshitzer Yu L, Mueller A H, Schiff D, Phys. Rev. C **58** (1998) 1706
- [15] Baier R, Schiff D, Zakharov B G, Ann. Rev. Nucl. Paer. Sci. **50** (2000) 37
- [16] Wang X N, Phys. Rev. Lett. **81** (1998) 2655, Phys. Rev. C **58** (1998) 2321
- [17] M. Gyulassy and M. Plumer, Nucl. Phys. A **527**, 641 (1991).
- [18] M. Gyulassy, M. Plumer, M. Thoma and X. N. Wang, Nucl. Phys. A **538**, 37C (1992) ; X. Wang and M. Gyulassy, Phys. Rev. Lett. **68**, 1480 (1992).

- [19] E. V. Shuryak, Phys. Rev. C **55**, 961 (1997) [arXiv:nucl-th/9605011].
- [20] M. G. Mustafa, D. Pal, D. K. Srivastava and M. Thoma, Phys. Lett. B **428**, 234 (1998) [arXiv:nucl-th/9711059].
- [21] Z. w. Lin and R. Vogt, Nucl. Phys. B **544**, 339 (1999) [arXiv:hep-ph/9808214]; Z. w. Lin, R. Vogt and X. N. Wang, Phys. Rev. C **57**, 899 (1998) [arXiv:nucl-th/9705006].
- [22] Y. L. Dokshitzer and D. E. Kharzeev, Phys. Lett. B **519**, 199 (2001) [arXiv:hep-ph/0106202].
- [23] K. Adcox *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **88**, 192303 (2002) [arXiv:nucl-ex/0202002].
- [24] Batsouli S, Kelly S, Gyulassy M, Nagle J L, Phys. Lett. B (2003) in press, nucl-th/0212068
- [25] X.-N. Wang, Phys. Rev. Lett. **81**, 2655 (1998).
- [26] M. Gyulassy and P. Levai, Phys. Lett. B **442**, 1 (1998) [arXiv:hep-ph/9807247]. ITATION = hep-ph 9804384;
- [27] Ter-Mikayelian M L, Dokl. Akad. Nauk SSSR **94** (1954) 1033
- [28] Ter-Mikayelian M L, High-Energy Electromagnetic Processes in Condensed Media, John Wiley & sons, New York (1972)
- [29] Kampfer B, Pavlenko O P, Phys. Lett. B **477** (2000) 171
- [30] Biro T S, Levai P, Muller B, Phys. Rev. D **42** (1990) 3078
- [31] Peshier A, Kampfer B, Pavlenko O P, Soff G, Phys. Rev. D **54** (1996) 2399
- [32] Levai P, Heinz U, Phys. Rev. C **57** (1998) 1879
- [33] O. K. Kalashnikov and V. V. Klimov, Sov. J. Nucl. Phys. **31**, 699 (1980).
- [34] V. V. Klimov, Sov. Phys. JETP **55** (1982) 199
- [35] R. D. Pisarski, PhysicaA **158** (1989) 146.
- [36] Rebhan A, Lect. Notes Phys. **583** (2002) 161, hep-ph/0105183; hep-ph/0111341.
- [37] Selikhov A V, Gyulassy M, Phys. Lett. B **316** (1993) 373; Phys. Rev. C **49** (1994) 1726.
- [38] Djordjevic M, Gyulassy M, Ter-Mikayelian effect for radiative energy loss in QCD matter, to be published
- [39] Djordjevic M, Gyulassy M, Radiative heavy quark energy loss in QCD matter, to be published
- [40] Gyulassy M, Levai P, Vitev I, Nucl. Phys. B **594** (2001) 371

- [41] Levai P, Papp G, Fai G, Gyulassy M, Barnafoldi G G, Vitev I, Zhang Y, Nucl. Phys. A **698** (2002) 631
- [42] E. Wang and X. N. Wang, Phys. Rev. Lett. **87**, 142301 (2001).